

# Important Billing and Coding Information for REBLOZYL® (luspatercept-aamt)

NDCs	
11-digit NDC: 59572-0711-01	REBLOZYL injection 25 mg/vial
11-digit NDC: 59572-0775-01	REBLOZYL injection 75 mg/vial

The red zero converts the 10-digit NDC to the 11-digit NDC. Payer requirements regarding the use of NDCs may vary. Electronic data exchange generally requires use of the 11-digit NDC.

HCPCS Codes	
J3490	Not otherwise classified drugs
J3590	Not otherwise classified biologics
J9999	Not otherwise classified anti-neoplastic drugs
C9399	Unclassified or biologicals (hospital outpatient use only)

CPT® Code	
96372	Therapeutic, prophylactic, or diagnostic injection (specify substance or drug); subcutaneous or intramuscular

How Supplied	
25-mg vial	25 mg lyophilized powder for solution for injection in a single-dose vial for reconstitution
75-mg vial	75 mg lyophilized powder for solution for injection in a single-dose vial for reconstitution

Billing Units	
1 mg = 1 unit	25-mg vial = 25 units 75-mg vial = 75 units

ICD-10-CM Diagnosis Code	
D56.1	<ul style="list-style-type: none"> <li>• Beta thalassemia major</li> <li>• Cooley's anemia</li> <li>• Homozygous beta thalassemia</li> <li>• Severe beta thalassemia</li> <li>• Thalassemia intermedia</li> <li>• Thalassemia major</li> </ul>
D56.5	<ul style="list-style-type: none"> <li>• Hemoglobin E-beta thalassemia</li> </ul>

Depending on payer preferences for billing and coding, the required miscellaneous J code for claim submission may vary. Therefore, payer preferences should be confirmed prior to submitting.

Note the use of a miscellaneous C code (C9399) for when REBLOZYL is used in a hospital infusion center.

**[Click here](#) for more information around billing and coding for REBLOZYL.**

The information contained herein is not intended to provide specific coding and reimbursement advice for any specific patient or situation. You should check with your coding specialist to ensure appropriate submissions.

Abbreviations: CPT, Current Procedural Terminology; HCPCS, Healthcare Common Procedure Coding System; ICD-10 CM, *International Classification of Diseases*, Tenth Revision, Clinical Modification; NDC, National Drug Code.

CPT® codes and descriptions are copyright 2019 American Medical Association (AMA). All rights reserved. CPT® is a registered trademark of the AMA.

**Please see Important Safety Information on the next page and full [Prescribing Information](#).**

**Reblozyl®**  
(luspatercept-aamt)  
for injection 25mg • 75mg

## INDICATION

**REBLOZYL is indicated for the treatment of anemia in adult patients with beta thalassemia who require regular red blood cell (RBC) transfusions.**

**REBLOZYL is not indicated for use as a substitute for RBC transfusions in patients who require immediate correction of anemia.**

## Important Safety Information

### WARNINGS AND PRECAUTIONS

#### Thrombosis/Thromboembolism

Thromboembolic events (TEE) were reported in 8/223 (3.6%) REBLOZYL-treated patients. TEEs included deep vein thrombosis, pulmonary embolus, portal vein thrombosis, and ischemic stroke. Patients with known risk factors for thromboembolism (splenectomy or concomitant use of hormone replacement therapy) may be at further increased risk of thromboembolic conditions. Consider thromboprophylaxis in patients at increased risk of TEE. Monitor patients for signs and symptoms of thromboembolic events and institute treatment promptly.

#### Hypertension

Hypertension was reported in 10.7% (61/571) of REBLOZYL-treated patients. Across clinical studies, the incidence of Grade 3 to 4 hypertension ranged from 1.8% to 8.6%. In patients with beta thalassemia with normal baseline blood pressure, 13 (6.2%) patients developed systolic blood pressure (SBP) >130 mm Hg and 33 (16.6%) patients developed diastolic blood pressure (DBP) >80 mm Hg. Monitor blood pressure prior to each administration. Manage new or exacerbations of preexisting hypertension using anti-hypertensive agents.

#### Embryo-Fetal Toxicity

REBLOZYL may cause fetal harm when administered to a pregnant woman. REBLOZYL caused increased post-implantation loss, decreased litter size, and an increased incidence of skeletal variations in pregnant rat and rabbit studies. Advise pregnant women of the potential risk to a fetus. Advise females of reproductive potential to use effective contraception during treatment and for at least 3 months after the last dose.

### ADVERSE REACTIONS

Serious adverse reactions occurring in 1% of patients included cerebrovascular accident and deep vein thrombosis. A fatal adverse reaction occurred in 1 patient treated with REBLOZYL who died due to an unconfirmed case of acute myeloid leukemia (AML).

Most common adverse reactions (at least 10% for REBLOZYL and 1% more than placebo) were headache (26% vs 24%), bone pain (20% vs 8%), arthralgia (19% vs 12%), fatigue (14% vs 13%), cough (14% vs 11%), abdominal pain (14% vs 12%), diarrhea (12% vs 10%) and dizziness (11% vs 5%).

### LACTATION

It is not known whether REBLOZYL is excreted into human milk or absorbed systemically after ingestion by a nursing infant. REBLOZYL was detected in milk of lactating rats. When a drug is present in animal milk, it is likely that the drug will be present in human milk. Because many drugs are excreted in human milk, and because of the unknown effects of REBLOZYL in infants, a decision should be made whether to discontinue nursing or to discontinue treatment. Because of the potential for serious adverse reactions in the breastfed child, breastfeeding is not recommended during treatment and for 3 months after the last dose.

**Please see full [Prescribing Information](#) for REBLOZYL.**

To find out more, call your Celgene Patient Support® Specialist at 1-800-931-8691, Monday–Friday, 8 AM–8 PM ET (*translation services available*).



REBLOZYL® and Celgene Patient Support® are registered trademarks of Celgene Corporation.

REBLOZYL® is licensed from Acceleron Pharma Inc.

All trademarks are the property of their respective owners.

© 2019 Celgene Corporation 11/19 US-RBZ-19-0074

**Reblozyl**®  
(luspaterecept-aamt)  
for injection 25mg • 75mg